

ISOMORPHISMS BETWEEN AFFINE HECKE ALGEBRAS

JIE-TAI YU

ABSTRACT. Let k be a field and suppose $p, q \in k$. We prove that the two affine Hecke algebras H_q and H_p of type A_n are isomorphic as k -algebras if and only if $p = q^{\pm 1}$.

There is a classical theorem that two finite dimensional Hecke algebras $\mathbb{H}_q$ and $\mathbb{H}_p$ over a field of characteristic 0 of same type are always isomorphic by Tits (see Bourbaki [B], see also Lusztig [L] for such an explicit isomorphism). Motivated by this, Xi [X] raised the question whether the same result holds for affine Hecke algebras and the question for the type A_2 was answered in [Y].

In this paper we answer the question of Xi for the type A_n for all $n > 2$. The methodology, involves the center (as a polynomial algebra) of the affine Hecke algebra and its automorphisms, has been adapted from Li and Yu[LY], where the question of isomorphism between quantum groups of type A is treated.

Let k be an arbitrary field and let H_q be the affine Hecke algebra of type $A_n (n \geq 2)$ over k with the generating set

$$\{T_i, X_j \mid i = 1, \dots, n; j = 1, \dots, n+1\}$$

subject to the following defining relations:

$$\begin{aligned} T_i T_i^{-1} &= 1 = T_i^{-1} T_i; \quad (T_i + 1)(T_i - q) = 0; \\ T_i T_{i+1} T_i &= T_{i+1} T_i T_{i+1}; \quad T_i T_j = T_j T_i (\forall \mid i - j \mid > 1); \\ X_j X_j^{-1} &= 1 = X_j^{-1} X_j; \quad X_i X_j = X_j X_i, \quad X_1 X_2 \dots X_{n+1} = 1; \\ T_i X_i T_i &= q X_i + 1; \quad X_j T_i = T_i X_j (\forall j \neq i; i+1) \end{aligned}$$

The center $Z(H_q)$ of H_q is

$$k[S_1, S_2, \dots, S_n],$$

2000 *Mathematics Subject Classification.* 16G10, 16S10, 16W20, 16Z05, 17B10, 20C30.

Key words and phrases. Affine Hecke groups, isomorphisms, automorphisms, center, polynomial algebras.

The research of Jie-Tai Yu was partially supported by an RGC-GRF grant.

where

$$S_i = S_i(X_1, \dots, X_n)$$

is the i -th elementary polynomial of $X_1, X_2, \dots, X_{n+1}$ (we omit X_{n+1} in the expression of S_i since $X_{i+1} = (X_1 \dots X_n)^{-1}$, so actually S_i are functions of $X_1, \dots, X_n$). Observe the symmetry that

$$S_i(X_1, \dots, X_n) = S_{n-i}(X_1^{-1}, \dots, X_n^{-1}).$$

Theorem 0.1. *Let H_q and H_p be isomorphic as k -algebras. Then $p = q^{\pm 1}$.*

Proof. Let H_q be isomorphic to H_p as k -algebras. Then we may assume $H_q = H_p$. In other words, there exist another generating set

$$\{t_i, x_j \mid i = 1, \dots, n; j = 1, \dots, n+1\}$$

subject to the following defining relations:

$$\begin{aligned} t_i t_i^{-1} &= 1 = t_i^{-1} t_i; \quad (t_i + 1)(t_i - q) = 0; \\ t_i t_{i+1} t_i &= t_{i+1} t_i t_{i+1}; \quad t_i t_j = t_j t_i \quad (\forall \mid i - j \mid > 1); \\ x_j x_j^{-1} &= 1 = x_j^{-1} x_j; \quad x_i x_j = x_j x_i, \quad x_1 x_2 \dots x_{n+1} = 1; \\ t_i x_i t_i &= q x_{i+1}; \quad x_j t_i = t_i x_j \quad (\forall j \neq i; i+1) \end{aligned}$$

The center $Z(H_q) = Z(H_p)$ of $H_q = H_p$ is

$$k[S_1, S_2, \dots, S_n] = k[s_1, s_2, \dots, s_n],$$

where $s_i = s_i(x_1, \dots, x_n)$ is the i -th elementary polynomial of

$$x_1, x_2, \dots, x_{n+1}.$$

Similarly

$$s_i(x_1, \dots, x_n) = s_{n-i}(x_1^{-1}, \dots, x_n^{-1}).$$

Obviously there exists a k -algebra isomorphism α from $k[X_1^{\pm 1}, \dots, X_n^{\pm 1}]$ onto $k[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$ taking X_j to x_j , and α induces an k -algebra automorphism of

$$Z(H_q) = k[S_1, S_2, \dots, S_n] = k[s_1, s_2, \dots, s_n] = Z(H_p)$$

taking S_i to s_i .

Obviously there exists an automorphism α of $Z(H_q)$ taking S_j to s_j for all j and α can only be lifted to the following isomorphisms from $k[X_1^{\pm 1}, \dots, X_n^{\pm 1}]$ onto $k[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$ taking X_i to $x_{\sigma(i)}$ for all i or taking X_i to $x_{\sigma(i)}^{-1}$ for some $\sigma \in \mathfrak{S}_n$. Note also $S_j(x_{\sigma(1)}, \dots, x_{\sigma(n)}) = s_j$ and $S_j(x_{\sigma(1)}^{-1}, \dots, x_{\sigma(n)}^{-1}) = s_{n-j}$ for all j and all $\sigma \in \mathfrak{S}_n$.

Now consider the scalar actions of the central elements

$$S_j, \quad s_j = S_j(x_{\sigma(1)}, \dots, x_{\sigma(n)})$$

(or $s_j = S_{n-j}(x_{\sigma(1)}^{-1}, \dots, x_{\sigma(n)}^{-1})$) in $Z(H) = Z(H_q) = Z(H_p)$ on the 1-dimensional simple $(H_q = H_p) = H$ -module M via the H_q -structure and H_p -structure of H , there must exist

$$a_1, a_2, \dots, a_n, a_{n+1} \in k^*, \quad a_1 a_2 \dots a_n a_{n+1} = 1$$

such that $a_i = q a_{i+1}$ for $i = 1, 2, \dots, n$ and

$$b_1, b_2, \dots, b_n, b_{n+1} \in k^*, \quad b_1 b_2 \dots b_n b_{n+1} = 1$$

such that $b_i = a_{\sigma(i)}$ for all i , or $b_i = a_{\sigma(i)}^{-1}$ for all i for some $\sigma \in \mathfrak{S}_n$ and $b_i = p b_{i+1}$ for all $i = 1, 2, \dots, n$, and

$$s_j(a_1, \dots, a_{n+1}) = s_j(b_1, \dots, b_{n+1})$$

for all j .

It is easy to see that the only possibilities are

- (1) $p = q$, $b_i = a_i$ for all i ; or
- (2) $p = q^{-1}$, $b_i = a_i^{-1}$ for all i .

□

Remark 2. By the methodology in this paper, we can treat the same isomorphism question for affine Hecke algebra of all other types and obtain the similar result.

Acknowledgements.

The author is grateful to Yangzhou University (Yangzhou), Yunnan Normal University (Kunming), Shanghai University (Shanghai), Osaka University and Kwansei Gakuin University (Osaka), Saitama University (Saitama), Chern Institute of Mathematical Research (Tianjin), Beijing International Center for Mathematical Research and Chinese Academy of Sciences (Beijing), and Academia Sinica (Taipei) for warm hospitality and stimulating environment during his visits in 2009 when this work was carry out.

REFERENCES

- [B] N. Bourbaki, *Lie groups and Lie algebras, Chapters 4-6*, Translated from the 1968 French original by Andrew Pressley. Elements of Mathematics (Berlin). Springer-Verlag, Berlin, 2002, ISBN: 3-540-42650-7.
- [L] G. Lusztig, *On a theorem of Benson and Curtis*, J. Algebra 71 (1981) 490-498.
- [LY] L.-B. Li and J.-T. Yu, *Isomorphisms between quantum groups $U_q(\mathfrak{sl}_{n+1})$ and $U_p(\mathfrak{sl}_{n+1})$* , arXiv:1001.0154.
- [X] N.-H. Xi, *Representations of affine Hecke algebras*, Lecture Notes in Mathematics, 1587, Springer-Verlag Berlin, 1994, ISBN: 3-540-58389-0.

- [Y] R. Yan, *Isomorphisms between two affine Hecke algebras of type $\widetilde{A}_2$* , Ph.D. Thesis, June 2009, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing.

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF HONG KONG, POKFU-LAM, HONG KONG SAR, CHINA

E-mail address: yujt@hkucc.hku.hk yujietai@yahoo.com